

CASE REPORT

Craniocerebral gunshot injury with no neurological deficit: the trending neurosurgical challenge.

Sharma Mitrajit¹, Barooah RK², Malakar Jaydeep³, Sudhy IK⁴

Received on May 10, 2019; editorial approval on July 20, 2019

ABSTRACT

Penetrating brain injury is a traumatic brain injury caused by high-velocity projectiles or low-velocity sharp objects. A wound in which the projectile breaches the cranium but does not exit is referred to as a penetrating wound. A large number of these patients who survive their initial wound will nevertheless expire shortly after admission to the hospital.¹ Here we discuss a case of penetrating bullet injury in a civilian, who presented with no neurological deficit and the management of the case along with a short discussion on the various presentation of the case and their respective neurosurgical management.

Keywords: Penetrating brain injury; civilian bullet injury; neurosurgery; neurology

INTRODUCTION

Penetrating brain injury (PBI) is a type of traumatic brain injury (TBI) defined as brain trauma in which a low-velocity sharp object, such as a knife, or a high-velocity projectile, such as a bullet, penetrates the skull but does not exit it.² Craniocerebral gunshot injuries (CGI) are currently increasingly encountered in civilian setting. Although less prevalent than closed head trauma, penetrating brain injury carries a grim prognosis.³ CGI are the most lethal of all firearm injuries, with reported survival rates of only 7% to 15%.⁴ This study reports an interesting case of CGI with no residual neurological defect.

CASE STUDY

A 14 year old male presented with a history of homicidal penetrating bullet injury to brain with facial soft tissue injury. The GCS (Glasgow Coma Scale) on presentation was E₄V₅M₆ with an entry wound seen at 4 cm above the right eyebrow with no exit wound. CT (computed tomography) showed a foreign body in the right parietal region with surrounding cerebral edema and hematoma formations in the track of the bullet (**Figure 1** and **2**). Triple intravenous antibiotics, anti epileptics and osmotic diuresis were started. He had no neurological or motor deficit.

Right parietal curvilinear incision deepened and temporalis muscle retracted. Craniotomy done a pellet of 4mm diameter is retrieved using bayonet forceps and continuous saline irrigation (**Figure 3**). Homeostasis achieved, duramater was not closed and bone flap repositioned. Post operative period was uneventful and case is in follow-up.

Address for correspondence:

¹Resident (**Corresponding author**)

Department of General Surgery

Mobile: +919706582392

Email: mitrajitsurgery@gmail.com

²Associate Professor

Department of Neurosurgery

^{3,4}Resident

Department of General Surgery

Gauhati Medical College and Hospital, Assam, India.

Cite this article as: Sharma Mitrajit, Barooah RK, Malakar Jaydeep, Sudhy IK. Craniocerebral gunshot injury with no neurological deficit: the trending neurosurgical challenge. *Int J Health Res Medico Leg Prae* 2020 January;6(1):67-70. DOI 10.31741/ijhrmlp.v6.i1.2020.14




Figure 1 CT scan showing the bullet position and bullet track




Figure 2 Scan showing the entry wound the bullet lodged in parietal lobe




Figure 3 The pellet in comparison to scale

DISCUSSION

The introduction of Guidelines for the Management of Penetrating Brain Injury has revolutionized the management of PBI during the last decade. There has been a paradigm shift toward a less aggressive debridement of deep seated fragments and a more aggressive antibiotic prophylaxis to improve outcomes.⁵ We started with ceftriaxone + tazobactam, amikacin and metronidazole. CT (brain) scanning is the neuroradiologic modality of choice. Guidelines for the Management of Penetrating Brain Injury recommends the treatment of small entrance bullet wounds to the head with no significant intracranial pathology, with local wound care and closure. More extensive wounds with nonviable scalp, bone, or dura should be debrided more extensively before primary closure or grafting to secure a watertight wound is to be done if possible within 12 hours of the injury. When the trajectory of the missile violates an open air sinus, a water tight closure of the dura should be done as the literature suggests that it may decrease the risk of abscess formation and CSF fistulas.⁶ Advanced age, suicide attempts, associated coagulopathy, Glasgow coma scale score of 3 with bilaterally fixed and dilated pupils, and high initial intracranial pressure have been correlated with worse outcomes in PBI patients

CONCLUSION

PBI, though less prevalent than closed head trauma, carries a worse prognosis. The publication of Guidelines for the

Management of Penetrating Brain Injury in 2001, attempted to standardize the management of PBI. A precise medical and surgical management of these inimitable injuries present a significant challenge. A better and standardized study has to be undertaken to bring in lime light the best modality of treatment of such cases in our infrastructure.

Consent: Written informed consent was obtained from the patient for publication of this case report and accompanying images.

Funding: None declared.

Conflict of interest: None declared.

Author disclosure: (1) The article is original with the author(s) and does not infringe any copyright or violate any other right of any third party. (2) The article has not been published (whole or in part) elsewhere, and is not being considered for publication elsewhere in any form, except as provided herein. (3) All author(s) have contributed sufficiently in the article to take public responsibility for it and (4) All author(s) have reviewed the final version of the above manuscript and approved it for publication.

REFERENCES

1. Sharma AP. Penetrating brain injury. *Journal of Lumbini Medical College*. 2013 Dec 30;1(2):112-7.
2. Modi M, Arivazhagan A, Bharath RD, Rao MB. Penetrating brain injury with machete, stuck to calvarium: Hurdles in imaging and solutions. *J of*

- neurosciences in rural practice. 2014 Nov;5(Suppl 1):S63.
3. Aarabi B, Alden TD, Chesnut R, Downs J, Ecklund J, Eisenberg H. Guidelines for the management of penetrating brain injury. *J Trauma*. 2001;51(2 Suppl):S1-86.
 4. Selden BS, Goodman JM, Cordell W, Rodman GH, Schnitzer PG. Outcome of self-inflicted gunshot wounds of the brain. *Annals of emergency medicine*. 1988 Mar 1;17(3):247-53.
 5. Nagib MG, Rockswold GL, Sherman RS, Lagaard MW. Civilian gunshot wounds to the brain: prognosis and management. *Neurosurgery*. 1986 May 1;18(5):533-7.
 6. Kazim SF, Shamim MS, Tahir MZ, Enam SA, Waheed S. Management of penetrating brain injury. *Journal of Emergencies, Trauma and Shock*. 2011 Jul;4(3):395.
 7. Prognosis in penetrating brain injury. *J Trauma*. 2001;51:S44-86.